

**Europäisches
Patentamt****European
Patent Office****Office européen
des brevets**

REC'D 21 OCT 2003

WIPO

PCT

Bescheinigung**Certificate****Attestation**

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

02078643.0

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk



Anmeldung Nr:
Application no.: 02078643.0
Demande no:

Anmeldetag:
Date of filing: 05.09.02
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Pantarhei Bioscience B.V.
P.O. Box 464
3700 AL Zeist
PAYS-BAS

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Pharmaceutical application of 15- or 16- substituted testosterone analogues

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)

Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

A61K31/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR

PHARMACEUTICAL APPLICATION OF 15- OR 16- SUBSTITUTED TESTOSTERONE
ANALOGUES

EPO - DG 1

5 TECHNICAL FIELD

05.09.2002

The present invention is in the field of steroid chemistry. More particularly, it relates to pharmaceutical dosage units containing 15- or 16-substituted testosterone analogues, as well as to therapeutic methods that employ these testosterone analogues. The testosterone analogues of the invention possess androgenic activity.

BACKGROUND OF THE INVENTION

At present, the main therapeutic application of androgenic steroids is in the treatment of hypogonadal males. Methods of treating androgen deficiency comprising the administration of androgens, such as testosterone, dihydrotestosterone, dehydroepiandrosterone and various esters of testosterone, or derivatives and analogues such as mesterolone are known in the art.

Three types of androgen deficiency in males are usually distinguished, i.e. primary androgen deficiency (testicular insufficiency), secondary androgen deficiency (hypothalamo-hypophyseal insufficiency) and androgen deficiency in ageing males (ADAM), also known as "male menopause" or "andropause".

As regards the long-term administration of androgens to males, a distinction can be made between therapy and supplementation. Therapy typically requires relatively high doses that are usually similar to the rate of production of endogenous androgens. Supplementation on the other hand is suitably done with dosages that are below the rate of production of endogenous androgens (i.e. testosterone, dihydrotestosterone and dehydroepiandrosterone).

Because of concerns about undesirable side effects, androgens are only used sparingly in both therapy and hormone supplementation. Indeed, androgens are normally only used for therapy in human males when primary or secondary androgen deficiency has been diagnosed.

Only a few androgens, e.g. dehydroepiandrosterone (DHEA) and 17 α -alkylated derivatives of testosterone, are suitable for oral administration because, unlike testosterone, they are largely resistant to hepatic metabolism. However, disadvantages of oral

administration are associated with the bad absorption of these androgens and the relatively high effect they exert on the liver and particular the liver metabolism (Bhasin et al. (1997) J. Clin. Endocr. Metab. 82:3-8). This is why, in existing protocols, androgens are generally administered in the form of 2-3 weekly depot injections or implants.

5 It has also been suggested in the prior art to use a combination of a progestogen and an androgen in a method of male contraception. In such a method the progestogen is administered in a sufficiently high amount to halt spermatogenesis - leading to azoospermia - and the androgen is co-administered to prevent androgen deficiency which would otherwise result from the administration of the progestogen. Because of concerns about reliability and
10 possible side-effects of the androgenic component, particularly when administered orally, male contraceptive have not yet made it beyond the experimental stage.

Other therapeutic uses of androgens that have been proposed in the prior art include treatment of the wasting syndrome and retro-viral drug induced lipodystrophia in HIV infected individuals, enhancement of recovery of critically ill catabolic individuals, treatment
15 of benign gynaecological disorders hormonal contraception in females, delayed puberty, female-to-male conversion.

As will be apparent from the above, there is an unmet need for androgens that (i) can be used effectively in the above mentioned therapeutic methods without causing undesirable side-effects, (ii) produce a very consistent, i.e. predictable, impact and (iii) may be
20 administered in a convenient manner.

SUMMARY OF THE INVENTION

25 The inventors have unexpectedly discovered that steroids which have not been used in therapeutic applications, i.e. 15-hydroxy or 16-hydroxy substituted testosterone analogues, meet the aforementioned requirements.

Surprisingly, it was found that the hydroxytestosterone analogues according to the present invention combine adequate androgenic potency with acceptable oral bioavailability
30 and minimum impact on liver metabolism. Consequently, the present androgens are particularly suited for any therapeutic applications for which androgens have been employed or recommended.

Another advantage of the hydroxytestosterone analogues according to the invention is their predictable and consistent therapeutic impact, which is believed to be related to the

efficiency of uptake and/or their low hepatic effect. In contrast, the therapeutic effect of known androgens that are sometimes administered orally, notably DHEA or 17 α -alkylated derivatives of testosterone, is very dependent on an individual's physiology and even his/her diet. Consequently, these androgens are usually applied in relatively high dosages in order to ensure that the minimum effective dosage is achieved in each individual. Naturally such high dosages have the disadvantage that they lead to relative overdosing in some individuals, which in turn is likely to produce pronounced side effects.

10 DETAILED DESCRIPTION OF THE INVENTION

Accordingly, one aspect of the invention relates to a pharmaceutical dosage unit containing at least 10 μ g of an androgenic steroid selected from the group consisting of 15-hydroxytestosterones, 16-hydroxytestosterones, precursors thereof and mixtures of these hydroxytestosterones and/or their precursors; and a pharmaceutically acceptable excipient.

The term "15-hydroxytestosterones" encompasses both 15 α -hydroxytestosterone (15 α ,17 β -dihydroxy-4-androsten-3-one) and 15 β -hydroxytestosterone (15 β ,17 β -dihydroxy-4-androsten-3-one). Similarly, the term "16-hydroxytestosterones" encompasses both 16 α -hydroxytestosterone (16 α ,17 β -dihydroxy-4-androsten-3-one) and 16 β -hydroxytestosterone (16 β ,17 β -dihydroxy-4-androsten-3-one).

The term "precursor" as used throughout this document refers to substances which are converted into one of the hydroxytestosterones of the invention following the administration of said precursor to a human subject.

The androgenic steroids of the present invention may suitably be administered by any enteral or parenteral methods known in the pharmaceutical art. The term "parenteral" as used in here encompasses transdermal, intravenous, intranasal, intravaginal, pulmonary, buccal, subcutaneous, intramuscular and intra-uterine modes of administration. The term "enteral" includes oral as well as rectal administration.

It has been shown that the hydroxytestosterones of the present invention occur in mammals, which means that they offer the important advantage that they can be labelled as natural (Yamazaki et al. Arch Biochem Biophys 1997 Oct 1;346(1):161-9). In addition, as can be derived from the aforementioned article, both 15 α -hydroxytestosterone, 15 β -hydroxytestosterone and 16 β -hydroxytestosterone occur in the human body, which means that *prima facie* the risk of undesired side-effects is lower than is the case for synthetic androgens

or natural androgens that are not found in the human body. Thus, in a particularly preferred embodiment of the present invention, the androgenic steroid is selected from the group consisting of 15α -hydroxytestosterone, 15β -hydroxytestosterone, 16β -hydroxytestosterone, precursors thereof and mixtures of these hydroxytestosterones and/or their precursors. Even
5 more preferably, the androgenic steroid is selected from the group consisting of 15α -hydroxytestosterone, 15β -hydroxytestosterone precursors thereof and mixtures of these hydroxytestosterones and/or their precursors.

The present dosage units may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such units may contain one or more
10 agents selected from the group consisting of sweetening agents, flavouring agents, colouring agents and preserving agents. Said dosage units may suitably contain a non-toxic pharmaceutically acceptable excipient. In case of e.g. tablets, these excipients may be, for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, such as maize starch,
15 or alginic acid; binding agents, such as starch, gelatine or acacia; and lubricating agents, such as magnesium stearate, stearic acid or talc. Tablets may be uncoated or may be coated by known techniques to delay disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate alone or with a wax may be employed.

20 Examples of dosage unit forms that may be administered by oral route include solutions, suspensions, emulsions, tablets, including sublingual and buccal tablets, soft gelatine capsules, including solutions used in soft gelatine capsules, aqueous or oil suspensions, emulsions, pills, lozenges, troches, tablets, syrups or elixirs and the like. Formulations for oral use may also be presented as hard gelatine capsules wherein the active
25 ingredient is mixed with an inert solid diluent, for example calcium carbonate, calcium phosphate or kaolin, or as soft gelatine capsules wherein the steroid component is mixed with water or an oil medium, such as peanut oil, liquid paraffin or olive oil.

Aqueous suspensions of the invention contain the steroids in a mixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients include a suspending
30 agent, such as sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia, and dispersing or wetting agents such as a naturally occurring phosphatide (e.g., lecithin), a condensation product of an alkylene oxide with a fatty acid (e.g., polyoxyethylene stearate), a condensation product of ethylene oxide with a long chain aliphatic alcohol (e.g.,

heptadecaethylene oxycetanol), a condensation product of ethylene oxide with a partial ester derived from a fatty acid and a hexitol (e.g., polyoxyethylene sorbitol mono-oleate), or a condensation product of ethylene oxide with a partial ester derived from fatty acid and a hexitol anhydride (e.g., polyoxyethylene sorbitan mono-oleate). The aqueous suspension may
5 also contain one or more preservatives such as ethyl or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents and one or more sweetening agents, such as sucrose, aspartame or saccharin. Ophthalmic formulations, as is known in the art, will be adjusted for osmotic pressure.

Oil suspensions may be formulated by suspending the steroid component in a
10 vegetable oil, such as arachide oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oil suspensions may contain a thickening agent, such as beeswax, hard paraffin or cetyl alcohol. Sweetening agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an antioxidant such as ascorbic acid.

15 Dispersible powders and granules of the invention suitable for preparation of an aqueous suspension by the addition of water may be formulated from the steroid component in a mixture with a dispersing, suspending and/or wetting agent, and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those disclosed above. Additional excipients, for example sweetening, flavouring and
20 colouring agents, may also be present.

The pharmaceutical dosage unit of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, such as olive oil or arachide oil, a mineral oil, such as liquid paraffin, or a mixture of these. Suitable emulsifying agents include
25 naturally-occurring gums, such as gum acacia and gum tragacanth, naturally occurring phosphatides, such as soybean lecithin, esters or partial esters derived from fatty acids and hexitol anhydrides, such as sorbitan mono-oleate, and condensation products of these partial esters with ethylene oxide, such as polyoxyethylene sorbitan mono-oleate. The emulsion may also contain sweetening and flavouring agents.

Syrups and elixirs may be formulated with sweetening agents, such as glycerol,
30 sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative, a flavouring or a colouring agent.

The pharmaceutical dosage units of the invention may also be in the form of a sterile injectable preparation, such as a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to the known art using those suitable dispersing or

wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, such as a solution of 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water and Ringer's solution, an isotonic sodium chloride. In addition, sterile fixed oils may conventionally be employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid may likewise be used in the preparation of injectables.

The steroids of this invention may also be administered in the form of suppositories for rectal administration of the drug. These dosage units can be prepared by mixing the steroid with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperatures and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and polyethylene glycols.

The present dosage unit preferably contains at least 20 μg , more preferably at least 40 μg and most preferably at least 60 μg of the androgenic steroid. In case the dosage unit is meant to provide a sustained release of the androgenic steroid, e.g. in case of a depot injection or a transdermal patch, the amount of androgenic steroid contained in the dosage unit may be as high as 1000 mg. Preferably the amount of the androgenic steroid does not exceed 500 mg, more preferably it does not exceed 200 mg. Most preferably the amount of androgenic steroid in the dosage unit does not exceed 100 mg.

In a particularly preferred embodiment of the invention, the dosage unit is designed for oral administration. Examples of oral dosage units which may suitably be used to deliver the present steroids include tablets, capsules, cachets, pellets, pills, powders or granules.

Typically the present pharmaceutical oral dosage unit has a weight in the range of 0.1-10 grams. The amount of androgenic steroid contained in the oral dosage unit is preferably at least 30 μg , more preferably at least 50 μg and most preferably at least 80 μg . Usually the amount of steroid contained in the oral dosage unit will not exceed 50 mg, more preferably it will not exceed 30 mg, most preferably it will not exceed 20 mg.

The hydroxytestosterone analogues of the present invention may advantageously be employed in a method of male contraception. As mentioned herein before, such a method requires the combined administration of a progestogen and an androgen. Typically the amount of progestogen included in the present dosage unit will exceed 10 μg , preferably it will exceed 30 μg . In a particularly preferred embodiment, the present dosage unit additionally contains a

progestogen in an amount equivalent to between 75 and 800 µg, more preferably between 100 and 500 µg levonorgestrel.

In other applications, e.g. in female hormone replacement therapy or female contraception, it may be advantageous to administer the present hydroxytestosterone analogues in combination with an estrogen. Accordingly, the present dosage unit, in addition to the present androgenic steroid, may suitably contain an estrogen, preferably in an amount of at least 10 µg, more preferably in an amount of at least 15 µg.

Examples of progestogens that may employed in the present dosage unit as well as in the methods described herein include progesterone, levonorgestrel, norgestimate,

norethisterone, dydrogesterone, drospirenone, 3-beta-hydroxydesogestrel, 3-keto desogestrel (=etonogestrel), 17-deacetyl norgestimate, 19-norprogesterone, acetoxypregnenolone, allylestrenol, anagestone, chlormadinone, cyproterone, demegestone, desogestrel, dienogest, dihydrogesterone, dimethisterone, ethisterone, ethynodiol diacetate, flurogestone acetate, gastrinon, gestodene, gestrinone, hydroxymethylprogesterone, hydroxyprogesterone, lynestrenol (=lynoestrenol), medrogestone, medroxyprogesterone, megestrol, melengestrol, nomegestrol, norethindrone (=norethisterone), norethynodrel, norgestrel (includes d-norgestrel and dl-norgestrel), norgestrienone, normethisterone, progesterone, quingestanol, (17alpha)-17-hydroxy-11-methylene-19-norpregna-4,15-diene-20-yn-3-one, tibolone, trimegestone, algestone acetophenide, nestorone, promegestone, 17-hydroxyprogesterone esters, 19-nor-17hydroxyprogesterone, 17alpha-ethinyl-testosterone, 17alpha-ethinyl-19-nor-testosterone, d-17beta-acetoxy-13beta-ethyl-17alpha-ethinyl-gon-4-en-3-one oxime, precursors of these compounds that are capable of liberating these progestogens *in vivo* when used in the present method and combinations thereof. Preferably the progestogen is selected from the group consisting of progesterone, desogestrel, etonogestrel, gestodene, dienogest, levonorgestrel, norgestimate, norethisterone, drospirenone, trimegestone, dydrogesterone, precursors of these progestogens and combinations thereof.

Examples of estrogens that may suitably be used in accordance with the present invention include ethinyl estradiol, mestranol, quinestranol, estradiol, estrone, estran, estriol, estetrol, conjugated equine estrogens, precursors thereof that are capable of releasing such an estrogen *in vivo* when used in the present method and combinations thereof. Preferably the estrogen is selected from the group consisting of ethinyl estradiol, estradiol, estetrol and combinations thereof.

Preferred precursors of the androgenic steroids according to the present invention are derivatives of the present hydroxytestosterones wherein the hydrogen atom of at least one

hydroxylgroup has been substituted by an acyl radical of a hydrocarbon carboxylic, sulfonic or sulfamic acid of 1-25 carbon atoms; tetrahydrofuranyl; tetrahydropyranal; or a straight or branched chain glycosydic residue containing 1-20 glycosidic units per residue.

Another aspect of the invention relates to a method of curatively or prophylactically treating a mammal, said method comprising the administration to said mammal of the present pharmaceutical dosage unit as described herein before. The present method is particularly suitable for treating humans.

Although an important aim of the present invention is to provide orally effective androgens, the present hydroxytestosterones and their precursors may suitably be administered by any enteral or parenteral routes known in the art. Preferably the mode of administration is selected from the group consisting of oral, transdermal, intranasal, intravaginal, intravenous, rectal, pulmonary, buccal, subcutaneous, intramuscular or intra-uterine administration. Even more preferably, the present method employs oral, intranasal, pulmonary, intravaginal or transdermal administration. Most preferably the method employs oral administration of the present androgenic steroid(s).

Oral, intravenous, rectal, intranasal, buccal and pulmonary administration are ideally suited for (at least) once daily administration. Transdermal administration is advantageously applied at frequencies between once a day and once a month. Intravaginal and intra-uterine administrations are advantageously operated at administration frequencies between once weekly and once monthly. Subcutaneous and intramuscular administration are suitably done in the form of depot injections at intervals of 1 week to 6 months, preferably at intervals of 4 weeks to 3 months.

For reasons of convenience and also to achieve high compliance rates, the present method preferably utilises administration intervals of 1 day, 1 week or 1 month. Regimens that employ once daily oral administration, once weekly transdermal administration or once monthly intravaginal or subcutaneous administration are particularly preferred. Most preferably the present method comprises at least once daily administration.

The androgenic steroid(s) of the invention are advantageously administered in an average daily amount which is at least 0.5 μg per kg of bodyweight. Preferably the average daily amount is at least 1 μg per kg of bodyweight, more preferably at least 1.5 μg per kg of bodyweight. The average daily amount usually does not exceed 1.5 mg per kg of bodyweight. Preferably the average daily amount does not exceed 1 mg per kg of bodyweight, more preferably it does not exceed 0.5 mg per kg of bodyweight.

The method according to the present invention is preferably employed in a method of treating or preventing androgen deficiency (in males or females); a method of hormonal contraception (in males or females); a method of treating or preventing wasting syndrome, anti-retroviral drug induced lipodystrophia, lack of well-being or fatigue in HIV infected individuals; a method of reversing catabolic state caused by a chronic illness, surgical intervention, oncological condition, trauma and/or malnutrition; a method of treating or preventing leydig cell dysfunction and germinal epithelial damage following cytotoxic chemotherapy; a method of treating or preventing fatigue or maintaining weight, hemoglobine or neutrophil count during or subsequent to cytotoxic chemotherapy or radiotherapy; a method of treating or preventing benign gynaecological disorders; a method of improving libido (in males or females); a method of treating or preventing delayed puberty; or a method of supporting female-to-male conversion. As mentioned herein before, the administration of the present steroids is particularly advantageous when employed in a method of treating or preventing androgen deficiency, a method of hormonal contraception or a method of treating or preventing benign gynaecological disorders. It is noted that symptoms of androgen deficiency can occur in both males and females. In a particularly preferred embodiment the present is used to treat or prevent androgen deficiency or employed as a contraceptive method in males.

The aforementioned methods usually employ continuous administration of the present steroids during a period of at least 10 days, preferably of at least 20 days. In a particular preferred embodiment the present steroids are administered during a period of at least 100 days with administration-free intervals that do not exceed 10 days, preferably do not exceed 8 days.

In a preferred embodiment, the present method comprises the co-administration of a progestogen in an average daily amount of at least 0.2 μg per kg of bodyweight and/or the an estrogen in an average daily amount of at least 0.2 μg per kg of bodyweight. Typically, the average daily amount of co-administered progestogen or estrogen will not exceed 1 mg per kg of bodyweight, more preferably said amount does not exceed 500 μg per kg of bodyweight, most preferably it does not exceed 300 μg per kg of bodyweight.

Yet another aspect of the invention relates to the use of an androgenic steroid selected from the group consisting of 15-hydroxytestosterones, 16-hydroxytestosterones, precursors thereof and mixtures of these hydroxytestosterones and/or their precursors in the preparation of a pharmaceutical composition, particularly a pharmaceutical dosage unit as described herein.

The invention is further illustrated by means of the following examples.

EXAMPLES

5

Example 1

An established competitive steroid binding assay was used to determine the relative binding affinity of 15α -hydroxytestosterone ($15\alpha,17\beta$ -dihydroxy-4-androsten-3-one), 15β -hydroxytestosterone ($15\beta,17\beta$ -dihydroxy-4-androsten-3-one) and 16β -hydroxytestosterone ($16\beta,17\beta$ -dihydroxy-4-androsten-3-one) to the androgen receptor (AR). Testosterone (17β -hydroxy-4-androsten-3-one) and 5α -dihydroxytestosterone (DHT, $5\alpha,17\beta$ -dihydroxy-androstan-3-one) served as controls in this assay.

The method employed was adapted from the scientific literature and described in detail by Chang et al (1987, J. Steroid Biochem., 27, 1-3, 123-131). Recombinant rat AR was expressed and purified from E. Coli. The *in vitro* assays involved the use of AR and [3 H]mibolerone, at a fixed concentration of 1.5 nM, as the labeled ligand. Recombinant AR was dissolved in binding buffer (50 mM Tris-Base, pH 7.5, 0.8 mM NaCl, 10% glycerol, 2 mM dithiothreitol, 1 mg/ml BSA and 2% ethanol) and duplicate aliquots were then incubated with [3 H]mibolerone at a final concentration of 1.5 nM, together with a vehicle control (1.0% DMSO), or the same amount of vehicle containing increasing concentrations of unlabeled steroid ligands as competitors. After incubation for 4 h at 4 $^{\circ}$ C, the unbound ligands were removed and the amounts of [3 H]mibolerone bound to the AR was measured. The average amounts of [3 H]mibolerone bound to AR at each concentration of competitor were used to make inhibition curves. IC₅₀ values were subsequently determined by a non-linear, least squares regression analysis. Inhibition constants (K_i) were calculated using the equation of Cheng and Prusoff (Cheng et al., 1973, Biochem. Pharmacol., 22, 3099-3108), using the measured IC₅₀ of the tested compounds, the concentration of radioligand employed in the assay, and the historical values for the K_d of the radioligand, which has been established as 3 nM.

Mean K_i values obtained for 15α -hydroxytestosterone, 15β -hydroxytestosterone, 16β -hydroxytestosterone, testosterone and DHT from three separate experiments are shown in Table 1. For comparison of binding affinities, the K_i value of DHT (which shows the highest affinity) was arbitrarily set at 100% and used to calculate the relative affinities (Table 1). As

compared to DHT and testosterone, 15 α -hydroxytestosterone, 15 β -hydroxytestosterone and 16 β -hydroxytestosterone bind with 2 to 3 orders of magnitude less affinity to AR (Table 1).

Table 1: Experimentally determined inhibition constants (K_i) for AR binding of 15 α -hydroxytestosterone, 15 β -hydroxytestosterone, 16 β -hydroxytestosterone, testosterone and 5 α -dihydroxytestosterone (DHT). Relative binding affinities are also shown, using K_i value of DHT as 100%.

Trivial Name (Chemical Name)	AR-binding	
	K _i (nM)	Relative affinity (%)
15 α -hydroxytestosterone (4-ANDROSTEN-15 α , 17 β -DIOL-3-ONE)	1820	0,05
15 β -hydroxytestosterone (4-ANDROSTEN-15 β , 17 β -DIOL-3-ONE)	557	0,18
16 β -hydroxytestosterone (4-ANDROSTEN-16 β , 17 β -DIOL-3-ONE)	814	0,12
Testosterone (4-ANDROSTEN-17 β -OL-3-ONE)	2	50
5 α -dihydroxytestosterone, DHT (5 α -ANDROSTAN-17 β -DIOL-3-ONE)	1	100

Example 2

To determine the *in vivo* androgenic potency of 15-substituted testosterone derivatives, 15 α -hydroxytestosterone (15 α ,17 β -dihydroxy-4-androsten-3-one) and 15 β -hydroxytestosterone (15 β ,17 β -dihydroxy-4-androsten-3-one) were administered orally, once daily, to immature male Wistar rats for 5 consecutive days. Testosterone (17 β -hydroxy-4-androsten-3-one) and 5 α -dihydroxytestosterone (DHT, 5 α ,17 β -dihydroxy-androstan-3-one), both administered orally, served as positive controls in this bioassay.

15 α -hydroxytestosterone, 15 β -hydroxytestosterone, testosterone and 5 α -dihydroxytestosterone were dissolved in an aqueous suspension of 2% Tween 80, at final test concentrations of 3 mg/ml and 1 mg/ml. Groups of 5 immature male Wistar rats, weighing 50 \pm 2 grams, received either vehicle treatment or were administered 30 or 10 mg/kg/day test substance by oral gavage in a dosing volume of 10 ml per kg for 5 consecutive days. During the experiment, animals were housed in cages of 45 x 23 x 15 cm and maintained in a controlled temperature of 22-24 oC and humidity (60-80%) environment with 12 hours light/dark cycles. Food (lab diet, rodent diet, PMI nutrition international) and water intake was *ad libitum*.

The animals were sacrificed 24 hours after receiving the final dose and the wet weight of the seminal vesicle of each animal was recorded and average wet weight per group was

calculated. Increase in average seminal vesicle wet weight relative to the average seminal vesicle wet weight of vehicle treated rats was considered indicative for *in vivo* androgenicity.

In vivo androgenicity data are shown in Table 2. Both 15 α and 15 β hydroxy-substituted analogues of testosterone induced a dose-dependent and significant increase in seminal vesicle wet weight. These *in vivo* androgenic effects were similar or equipotent to the dose-dependent effects observed for testosterone and considerably more potent than the *in vivo* androgenicity observed after oral administration of DHT, a classical androgen and the active metabolite of testosterone *in vivo* (O'Donnel et al., 1996, Endocrinology, 137, 2703-2710).

Table 2: Average increase in seminal vesicle wet weight relative to the average wet weight of vehicle (2% Tween-80) treated animals for groups of 5 immature Wistar rats, treated once daily, orally, for five consecutive days with 15 α -hydroxytestosterone, 15 β -hydroxytestosterone, testosterone or 5 α -dihydroxytestosterone (DHT). Relative androgenic potencies are also shown, using DHT as reference (100%).

Trivial Name (Chemical Name)	In vivo androgenicity		
	Oral dose mg/kg/day	Increase seminal vesicle wet weight (%)	Relative potency* (%)
15 α -hydroxytestosterone (4-ANDROSTEN-15 α , 17 β -DIOL-3-ONE)	30	40	148
	10	28	560
15 β -hydroxytestosterone (4-ANDROSTEN-15 α , 17 β -DIOL-3-ONE)	30	35	130
	10	20	400
Testosterone (4-ANDROSTEN-17 β -OL-3-ONE)	30	47	174
	10	41	820
5 α -dihydroxytestosterone, DHT (5 α -ANDROSTAN-17 β -DIOL-3-ONE)	30	27	100
	10	5	100

05.09.2002

CLAIMS

(95)

1. A pharmaceutical dosage unit containing at least 10 µg of an androgenic steroid selected from the group consisting of 15-hydroxytestosterones, 16-hydroxytestosterones,
5 precursors thereof and mixtures of these hydroxytestosterones and/or their precursors; and a pharmaceutically acceptable excipient.
2. The pharmaceutical dosage unit according to claim 1, wherein the steroid is selected from the group consisting of 15α-hydroxytestosterone, 15β-hydroxytestosterone, 16β-
10 hydroxytestosterone, precursors thereof and mixtures of these hydroxytestosterones and/or their precursors.
3. The pharmaceutical dosage unit according to claim 2, wherein the steroid is selected from the group consisting of 15α-hydroxytestosterone, 15β-hydroxytestosterone, precursors
15 thereof and mixtures of these hydroxytestosterones and/or these precursors.
4. The pharmaceutical dosage unit according to claim 3, wherein the steroid is selected from the group consisting of 15α-hydroxytestosterone, precursors thereof and mixtures of 15α-
20 hydroxytestosterone and/or these precursors.
5. The pharmaceutical dosage unit according to claim 3, wherein the steroid is selected from the group consisting of 15β-hydroxytestosterone, precursors thereof and mixtures of 15β-
hydroxytestosterone and/or these precursors.
- 25 6. The pharmaceutical dosage unit according to claim 2, wherein the steroid is selected from the group consisting of 16β-hydroxytestosterone, precursors thereof and mixtures of 16β-
hydroxytestosterone and/or these precursors.
7. The pharmaceutical dosage unit according to any one of claims 1-6, wherein the dosage
30 unit is designed for oral administration.
8. The pharmaceutical dosage unit according to claim 7, wherein the dosage unit is a tablet, a capsule, a cachet, a pellet, a pill, a powder or granules.

9. The pharmaceutical dosage unit according to any one of claims 1-7, wherein the dosage unit contains between 20 μ g and 1000 mg, preferably between 40 μ g and 500 mg of the androgenic steroid.
- 5 10. The pharmaceutical dosage unit according to any one of claims 1-8, wherein the dosage unit additionally contains at least 10 μ g of a progestogen and/or at least 10 μ g of an estrogen.
- 10 11. The pharmaceutical dosage unit according to any one of claims 1-10 for use in a method of curatively or prophylactically treating a mammal, said method comprising the administration of the dosage unit to said mammal.
12. The pharmaceutical dosage unit according to claim 11, wherein the method comprises oral administration.
- 15 13. The pharmaceutical dosage unit according to claim 11 or 12, wherein the method comprises the administration of the steroid in an average daily amount in the range of 0.5 μ g to 1.5 mg per kg of bodyweight, preferably of 1 μ g to 1 mg per kg of bodyweight.
- 20 14. The pharmaceutical dosage unit according to any one of claims 11-13 for use in a method of treating or preventing androgen deficiency; a method of hormonal contraception; a method of treating or preventing wasting syndrome, anti-retroviral drug induced lipodystrophia, lack of well-being or fatigue in HIV infected individuals; a method of reversing catabolic state caused by a chronic illness, surgical intervention, oncological condition, trauma and/or malnutrition; a method of treating or preventing leydig cell
25 dysfunction and germinal epithelial damage following cytotoxic chemotherapy; a method of treating or preventing fatigue or maintaining weight, hemoglobine or neutrophil count during or subsequent to cytotoxic chemotherapy or radiotherapy; a method of treating or preventing benign gynaecological disorders; a method of improving libido; a method of
30 treating or preventing delayed puberty; or a method of supporting female-to-male conversion.
15. Use of an androgenic steroid selected from the group consisting of 15-hydroxytestosterones, 16-hydroxytestosterones, precursors thereof and mixtures of these

hydroxytestosterones and/or their precursors in the preparation of a pharmaceutical composition.

05.09.2002

ABSTRACT

95

The present invention relates to pharmaceutical dosage units containing 15- or 16-substituted testosterone analogues, as well as to therapeutic methods that employ these testosterone analogues. More particularly, the present invention is concerned with a pharmaceutical dosage unit containing at least 10 µg of an androgenic steroid selected from the group consisting of 15-hydroxytestosterones, 16-hydroxytestosterones, precursors thereof and mixtures of these hydroxytestosterones and/or their precursors; and a pharmaceutically acceptable excipient.

The term "15-hydroxytestosterones" encompasses both 15 α -hydroxytestosterone (15 α , 17 β -dihydroxy-4-androsten-3-one) and 15 β -hydroxytestosterone (15 β , 17 β -dihydroxy-4-androsten-3-one). Similarly, the term "16-hydroxytestosterones" encompasses both 16 α -hydroxytestosterone (16 α , 17 β -dihydroxy-4-androsten-3-one) and 16 β -hydroxytestosterone (16 β , 17 β -dihydroxy-4-androsten-3-one).

The androgenic steroids according to the invention are advantageously employed in e.g. a method of treating or preventing androgen deficiency or a method of hormonal contraception.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☒ **FADED TEXT OR DRAWING**

☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.